

ANNALES ZOOLOGICI FENNICI

Vol. 14 • 1977

PUBLISHED BY

SOCIETAS BIOLOGICA FENNICA VANAMO

SNELLMANINKATU 9 — 11, SF-00170 HELSINKI 17, FINLAND

Correction

Owing to an editorial error the issue No. 3 begins with page 119. Pages 119—124 were, however, already included in No. 2. Therefore the pages 119—124 occur twice in this volume.

Oikaisu

Tämän niteen kolmannen numeron ensimmäinen sivu on 119. Sivut 119—124 sisältyivät kuitenkin jo numeroon 2, joten nämä sivunumerot ovat tässä niteessä toimituksen tekemän erehdyksen takia kahteen kertaan.

Contents — Sisältö

HALKKA, O., RAATIKAINEN, M., HALKKA, L. & RAATIKAINEN, T.: Coexistence of four species of spittle-producing Homoptera	228—231
HALLAS, T. E.: Survey of the Tardigrada of Finland	173—183
HAUKIOJA, E. & NIEMELÄ, P.: Retarded growth of a geometrid larva after mechanical damage to leaves of its host tree	48— 52
HELLE, P., HYYTINEN, L. & ITÄMIES, J.: Records of bird-flies (Diptera, Hippoboscidae) from northern Finland	232—234
HOKKANEN, H., TÖRMÄLÄ, T. & VUORINEN, H.: Seasonal changes in the circadian activity of <i>Pteromys volans</i> L. in central Finland	94— 97
HOLOPAINEN, I. & PAASIVIRTA, L.: Abundance and biomass of the meiozoobenthos in the oligotrophic and mesohumic lake Pääjärvi, southern Finland	124—134
ITÄMIES, J., & OJANEN, M.: Autumn predation of <i>Parus major</i> and <i>P. montanus</i> upon two leaf-mining species of <i>Lithocolletis</i> (Lepidoptera, Lithocolletidae)	235—241
JANSSON, A.: Distribution of Micronectae (Heteroptera, Corixidae) in Lake Päijänne, central Finland: Correlation with eutrophica- tion and pollution	105—117
JANSSON, A.: Micronectae (Heteroptera, Corixidae) as indicators of water quality in two lakes in southern Finland	118—124
KOSKELA, H. & HANSKI, I.: Structure and succession in a beetle community inhabiting cow dung	204—223
NURMINEN, M.: Enchytraeidae (Oligochaeta) from the Gross- glockner region of the Austrian Alps	224—227
OIKARI, A.: Renal adaptations of Baltic <i>Myoxocephalus scorpius</i> (L.) and <i>M. quadricornis</i> (L.) to hypotonic brackish water	162—182
PAJUNEN, V. I.: Population structure in rock-pool Corixids (Hemiptera, Corixidae) during the reproduction season	26— 47
SARVALA, J.: The naupliar development of six species of fresh- water harpacticoid Copepoda	135—161
SARVALA, J. & RANTA, E.: Performance of the Muus sampler in surveys of the brackish-water macrofauna	191—197
SISULA, H. & VIRTANEN, E.: Effects of storage on the energy and ash content of biological material	119—123
TÄHTI, H. & SOIVIO, A.: Respiratory and circulatory differences between induced and spontaneous arousals in hibernating hedgehogs (<i>Erinaceus europaeus</i> L.)	198—213

TURUNEN, S.: Changes in soluble proteins and esterases during embryogenesis in <i>Pieris brassicae</i> (L.) (Lepidoptera, Pieridae)	185—190
UTRIO, P. & ERIKSSON, K.: Volatile fermentation products as attractants for Macrolepidoptera	98—104
VÄISÄNEN, R. A.: Geographic variation in timing of breeding and egg size in eight European species of waders	1— 25
VIITALA, J.: Social organization in cyclic subarctic populations of the voles <i>Clethrionomys rufocanus</i> (Sund.) and <i>Microtus</i> <i>agrestis</i> (L)	53— 93